

Tonsillectomy (adult)

ENT04 Lite - Expires end of January 2024



This fact sheet is for general information about this procedure only. It is not intended to be used as medical advice or to replace advice that your relevant healthcare professional would give you. If you have a particular medical problem, please consult a healthcare professional.

© EIDO Systems International Limited. The operation and treatment information in this document is published under license from EIDO Systems International and is protected by copyright laws. Other than for your personal, non-commercial use, you may not copy, print out, download or otherwise reproduce any of the information.

eidohealthcare.com



What is a tonsillectomy?

A tonsillectomy is an operation to remove the tonsils. The tonsils are part of a group of lymphoid tissues (like the glands in your neck) that help to fight off infection.

What are the benefits of surgery?

Tonsillitis happens if the tonsils become infected. This causes pain, fever and difficulty swallowing and can make you feel unwell.

Your doctor may be concerned that there is an underlying cancer, especially if one of your tonsils is larger than the other, has an ulcer on it or you have confirmed cancer in the glands of your neck.

Are there any alternatives to surgery?

Surgery is recommended as it is the only dependable way to stop tonsillitis that keeps on coming back.

In children, a long course of antibiotics may break a cycle of frequent infections or the tonsillitis may simply stop after a few years. For adults, this treatment is less likely to be effective, especially following glandular fever.

What will happen if I decide not to have the operation or the operation is delayed?

You will probably keep on getting tonsillitis.

Untreated tonsillitis can sometimes form an abscess behind the tonsil. This is known as a 'quinsy'. The pus will need to be removed from the abscess through a needle, and you will need antibiotics.

Rarely, the infection may spread further into the tissues of your neck, causing a 'parapharyngeal' or 'retropharyngeal' abscess. This is a serious complication and needs an operation to drain away the pus.

If your tonsillitis does not get better with your usual treatment, contact your healthcare team. This may be a sign that the infection has spread further into the tissues of your neck, causing a 'parapharyngeal' or 'retropharyngeal' abscess.

What does the operation involve?

The operation is performed under a general anaesthetic and usually takes about 30 minutes.

Your surgeon will perform the tonsillectomy through your mouth using one of the following techniques.

- Cold dissection technique – Your surgeon will use a steel instrument to peel or cut the tonsil away from the layer of muscle underneath it
- Diathermy technique – Your surgeon will use a special instrument that uses heat to stop the bleeding

How can I prepare myself for the operation?

If you smoke, stopping smoking now may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.


Speak to the healthcare team about any vaccinations you might need to reduce your risk of serious illness while you recover. When you come into hospital, practise social distancing and hand washing and wear a face covering when asked.

What complications can happen?

Some complications can be serious and can even cause death.

General complications of any operation

- Bleeding on the day of surgery
- Infection of the surgical site (wound)
- Allergic reaction to the equipment, materials or medication
- Blood clot in your leg
- Blood clot in your lung



Tonsillitis

- Chest infection

Specific complications of this operation

- Small pieces of the tonsil may be left behind
- Lingual tonsillitis
- Change of taste
- Feeling you have something in your throat
- Change in your voice

Consequences of this procedure

- Pain

How soon will I recover?

You should be able to go home the next day.

The pain can last for up to 2 weeks. It usually gets gradually worse over the first 4 to 5 days before beginning to improve, and tends to be worse first thing in the morning.

You will need to stay off work or school and away from groups of people for 2 weeks. This is to help prevent throat infections while your throat is still healing.

Regular exercise should help you to return to normal activities as soon as possible. Before you

start exercising, ask the healthcare team or your GP for advice.

Most people make a full recovery and return to normal activities.

Summary

Surgery is the only dependable way to stop tonsillitis that keeps on coming back and the resulting pain, fever and difficulty swallowing.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewer

Matthew Campbell (MBBS, FRACS)

Illustrator

Medical Illustration Copyright © Nucleus Medical
Art. All rights reserved. www.nucleusinc.com